arXiv:1612.06268v2 [math.NT] 31 Dec 2016

Product formulas for the 5-division points on the Tate normal form E_5.

Patrick Morton

December 17, 2016

Abstract

Explicit formulas are proved for the 5-torsion points on the Tate normal form E_5 of an elliptic curve having $(X,Y) = (0,0)$ as a point of order 5. These formulas express the coordinates of points in $E_5[5] - \langle(0,0)\rangle$ as products of linear fractional quantities in terms of 5-th roots of unity and a parameter u, where the parameter b which defines the curve E_5 is given as $b = (\varepsilon^5 u^5 - \varepsilon^{-5})/(u^5+1)$ and $\varepsilon = (-1+\sqrt{5})/2$.

1 Introduction.

In previous papers, several new formulas for the 3-division points on the Deuring normal form

$$E_3 :\ Y^2 + \alpha XY + Y = X^3,$$

and the 4-division points on the Tate normal form

$$E_4 :\ Y^2 + XY + bY = X^3 + bX^2,$$

have recently been given. For the curve E_3, the point

$$(X,Y) = \left(\frac{-3\beta}{\alpha(\beta-3)}, \frac{\beta - 3\omega}{\beta - 3}\right)$$

represents the six points of order 3 in $E_3[3] - \langle(0,0)\rangle$, where ω is one of the two primitive cube roots of unity and (α,β) lies on the Fermat cubic

$$Fer_3 :\ 27X^3 + 27Y^3 = X^3Y^3.$$

(See [5].) Setting $b = 1/\alpha^4$ in the equation for E_4, a point of order 4 in $E_4[4] - \langle(0,0)\rangle$ is the point

$$(X, Y) = \left(-\beta_1\beta_2\beta_3, \beta_1^2\beta_2^2\beta_3\right), \tag{1.1}$$

where

$$\beta_n = \frac{\beta + 2i^n}{2\beta}, \quad i = \sqrt{-1},$$

and the point (α, β) lies on the Fermat quartic

$$Fer_4 : \ 16X^4 + 16Y^4 = X^4Y^4.$$

Replacing β by $i\beta$ (so β_n becomes β_{n-1}) and i by $-i$ in the above formula yields 8 of the 12 points of order 4 in $E_4[4]$. The other points of order 4 are $(0,0), (0,-b) \in \langle(0,0)\rangle$ and the two points $(-2b, 2\beta_1\beta_3 b)$ and $(-2b, 2\beta_2\beta_4 b)$. (See [2], [3].)

Similar formulas have been given for the 6-torsion points on the Tate normal form E_6, by Lynch [2]. This normal form is

$$E_6 : \ Y^2 + aXY + bY = X^3 + bX^2, \quad b = -(a-1)(a-2).$$

Lynch's formulas express the coordinates of 6-torsion points on E_6 as products of linear fractional quantities in α and β and a cube root of unity ω, where (α, β) is a point on the elliptic curve

$$Y^2 = X^3 + 1$$

and the parameter a is given by

$$a = \frac{10\beta^2 - 18}{9(\beta^2 - 1)} = \frac{10\alpha^3 - 8}{9\alpha^3}.$$

The exact formulas are somewhat complicated; these can be found in [2].

In this note I will prove analogous formulas for the points of order 5 on the Tate normal form

$$E_5(b) : \ Y^2 + (1+b)XY + bY = X^3 + bX^2,$$

on which the point $(0,0)$ is a point of order 5. (See the discussion in [4] for more on the Tate normal form.) These formulas are similar to the expressions

(1.1) for the points of order 4 on the curve E_4, in that they are products of linear fractional quantities in a parameter u, where

$$b = \frac{\varepsilon^5 u^5 + \bar{\varepsilon}^5}{u^5 + 1};$$

here $\varepsilon = \frac{-1+\sqrt{5}}{2}$ and $\bar{\varepsilon} = \frac{-1-\sqrt{5}}{2}$ is its algebraic conjugate. These expressions are quite a bit simpler than the formulas given by Verdure [9], in which the Y-coordinates of the 5-torsion points are expressed in terms of a formal root x_0 of the 5-division polynomial. In this paper, the quantity

$$u^5 = -\frac{b - \bar{\varepsilon}^5}{b - \varepsilon^5},$$

which is up to sign the same as Verdure's Kummer element (see Theorem 5 in [9]), arises naturally in the process of solving the quintic equation $g(X) = 0$ below using Watson's method (see [1] and Section 4 of this paper).

The expressions given here (see Theorem 2.1) also allow one to check "by hand" that these points do indeed have order 5, assuming that they represent points on E_5. (See Theorem 3.1 and the discussion in Section 3.) These formulas will be used in a forthcoming paper to prove the case $p = 5$ of the conjectures stated in [6] and [7].

2 Points of order 5 on $E_5(b)$.

The X-coordinates of points of order 5 on $E_5(b)$ which are not in the group

$$\langle (0,0) \rangle = \{O, (0,0), (0,-b), (-b,0), (-b,b^2)\}$$

are roots of the polynomial

$$\begin{aligned} D_5(x) = {} & 5x^{10} + (5 + 25b + 5b^2)x^9 + (1 + 38b + 44b^2 + 7b^3 + b^4)x^8 \\ & + (9b + 127b^2 + 26b^3 + 3b^4 - b^5)x^7 + (36b^2 + 248b^3 + 19b^4 - 3b^5 + b^6)x^6 \\ & + (84b^3 + 322b^4 + 71b^5 + 3b^6 - b^7)x^5 + (126b^4 + 293b^5 + 94b^6 + 12b^7 + b^8)x^4 \\ & + (125b^5 + 180b^6 + 50b^7 + 5b^8)x^3 + (80b^6 + 65b^7 + 10b^8)x^2 \\ & + (30b^7 + 10b^8)x + 5b^8. \end{aligned}$$

This follows easily from [8], Exercise 3.7 (p. 105), applied to the curve E_5, after factoring out $x(x + b)$ from the polynomial $\psi_5(x)$. (But note that the

formula for b_2 on p. 42 should be $b_2 = a_1^2 + 4a_2$.) This polynomial factors into 5 times the product of two polynomials

$$\begin{aligned} g(X) &= X^5 + \frac{1}{20}(\alpha - 5)(-3 - \alpha - 7b + 3b\alpha - 2b^2)X^4 \\ &\quad + \frac{\alpha}{5}b(1 + 2\alpha - 11b + 4b\alpha - b^2)X^3 \\ &\quad + \frac{1}{10}(\alpha - 5)b^2(-9 - 2\alpha - 6b + b\alpha - b^2)X^2 + (3b^3 + b^4)X + b^4, \end{aligned}$$

where $\alpha^2 = 5$. Using Watson's method of solving the for the roots of a quintic equation from [1], we find that the roots of $g(X)$ are given by

$$\begin{aligned} X &= \frac{(5-\alpha)}{100}\{(-18 + 8\alpha - 12b + 6b\alpha - 2b^2)u^4 + (-7 + 3\alpha + 12b - 4b\alpha + 2b^2)u^3 \\ &\quad + (-3 + \alpha - 7b + 7b\alpha - 2b^2)u^2 + (-2 + 22b + 2b^2)u - 3 - \alpha - 7b + 3b\alpha - 2b^2\} \\ &= \frac{(5-\alpha)}{100}(A_4u^4 + A_3u^3 + A_2u^2 + A_1u + A_0), \end{aligned}$$

where

$$u^5 = \phi(b) = \frac{2b + 11 + 5\alpha}{-2b - 11 + 5\alpha} = \frac{b - \bar{\varepsilon}^5}{-b + \varepsilon^5},$$
$$\varepsilon = \frac{-1 + \alpha}{2} = \zeta + \zeta^4, \quad \bar{\varepsilon} = \frac{-1 - \alpha}{2} = \zeta^2 + \zeta^3,$$

and ζ is a primitive 5-th root of unity. (The details of Watson's method applied to the polynomial $g(X)$ are given in the Appendix. Note that ε and $\bar{\varepsilon}$ are the quadratic Gaussian periods for $\mathbb{Q}(\zeta)$.) This may be verified on Maple by plugging the expression for X into $g(X)$, and using the formula

$$b = \frac{\varepsilon^5 u^5 + \bar{\varepsilon}^5}{u^5 + 1} \tag{2.1}$$

for b in terms of u.

Using this formula for b, the above value of X can also be written as

$$X = \frac{(-7 + 3\alpha)}{4}\frac{(-2u^2 + (1+\alpha)u - 3\alpha - 7)(2u^2 + (4 + 2\alpha)u + 3\alpha + 7)}{(-2u^2 + (1+\alpha)u - 2)(u+1)^2}. \tag{2.2}$$

The formulas (2.1) and (2.2) show that there are 10 such values, since replacing u by $\zeta^i u$ (and leaving α unchanged), or replacing α by $-\alpha$ and u by

$1/(\zeta^i u)$ gives the other X-coordinates. Setting $\alpha = \zeta - \zeta^2 - \zeta^3 + \zeta^4$, this expression factors:

$$X = -\varepsilon^4 \frac{[u-(1+\zeta)^2][u-\zeta(1+\zeta)^2][u-\zeta^2(1+\zeta)^2][u-\zeta^3(1+\zeta)^2]}{(u+\zeta^2)(u+\zeta^3)(u+1)^2}.$$

The zeros and poles of this function of u are all units in $\mathbb{Q}(\zeta_5)$. We will now show that the corresponding Y-coordinates factor in a similar way. We derive the following theorem using calculations in an extension of the field $\mathbb{Q}(\zeta, b)$, but the formulas themselves are valid over any field whose characteristic is different from 5.

Theorem 2.1. *If $b = \frac{\varepsilon^5 u^5 + \bar{\varepsilon}^5}{u^5+1}$, the X-coordinates of the points of order 5 in $E_5[5] - \langle(0,0)\rangle$ are given by the formula*

$$X = -\varepsilon^4 \frac{[u-(1+\zeta)^2][u-\zeta(1+\zeta)^2][u-\zeta^2(1+\zeta)^2][u-\zeta^3(1+\zeta)^2]}{(u+\zeta^2)(u+\zeta^3)(u+1)^2}, \quad (2.3)$$

where $\varepsilon = \frac{-1+\alpha}{2}$, $\alpha = \pm\sqrt{5} = \zeta - \zeta^2 - \zeta^3 + \zeta^4$, and $\zeta = \zeta_5$ is a primitive 5-th root of unity. The corresponding Y-coordinates are given by

$$Y_1 = \varepsilon^7 \frac{[u-(1+\zeta)^2]^2[u-\zeta(1+\zeta)^2]^2[u-\zeta^2(1+\zeta)^2]^2[u-\zeta^3(1+\zeta)^2]}{(u+\zeta^2)^2(u+\zeta^3)(u+\zeta^4)(u+1)^3},$$

and

$$Y_2 = \varepsilon^7 \frac{[u-(1+\zeta)^2][u-\zeta(1+\zeta)^2]^2[u-\zeta^2(1+\zeta)^2]^2[u-\zeta^3(1+\zeta)^2]^2}{(u+\zeta)(u+\zeta^2)(u+\zeta^3)^2(u+1)^3}.$$

Proof. Putting (2.1) and (2.2) into the equation for E_5 yields the following equation for Y:

$$AY^2 + BY + C = 0,$$

where

$$\begin{aligned}
A =& \frac{1}{8}(2u^2 + (-1+\alpha)u + 2)(-2u^2 + (1+\alpha)u - 2)^3(u+1)^6;\\
B =& \frac{-\varepsilon^7}{16}(-4u^3 + (3+\alpha)u^2 - 2(1+\alpha)u + 6 + 2\alpha)(-2u^2 + (1+\alpha)u - 7 - 3\alpha)\\
&\times(-2u^2 + (1+\alpha)u - 2)(2u^2 + (4+2\alpha)u + 7 + 3\alpha)^2(u+1)^3;\\
C =& \frac{\varepsilon^{14}}{64}(-2u^2 + (1+\alpha)u - 7 - 3\alpha)^3(2u^2 + (4+2\alpha)u + 7 + 3\alpha)^4.
\end{aligned}$$

The discriminant of the quadratic is

$$\begin{aligned} D =& \frac{-5\alpha\varepsilon^{13}}{64} u^4(-2u^2+(1+\alpha)u-7-3\alpha)^2(-2u^2+(1+\alpha)u-2)^2 \\ & \times(2u^2+(4+2\alpha)u+7+3\alpha)^4(u+1)^6, \end{aligned}$$

which is $-\alpha\varepsilon = (\zeta^2-\zeta^3)^2$ times a square. Thus, the roots of the quadratic are

$$Y = \frac{-B \pm (\zeta^2-\zeta^3)\alpha\frac{\varepsilon^6}{8}S}{2A} = \frac{-16B \pm 2(\zeta^2-\zeta^3)(\zeta-\zeta^2-\zeta^3+\zeta^4)\varepsilon^6 S}{32A},$$

where

$$S = u^2(-2u^2+(1+\alpha)u-7-3\alpha)(-2u^2+(1+\alpha)u-2)(2u^2+(4+2\alpha)u+7+3\alpha)^2(u+1)^3.$$

Now, $1/\varepsilon = -\bar{\varepsilon} = -(\zeta^2+\zeta^3)$, which gives that

$$(\zeta^2-\zeta^3)(\zeta-\zeta^2-\zeta^3+\zeta^4)\varepsilon^6 = \varepsilon^7(-\zeta^3+3\zeta^2+2\zeta+1).$$

The numerator in the expression for Y then becomes

$$\begin{aligned} -16B \pm 2(-\zeta^3+3\zeta^2+2\zeta+1)\varepsilon^7 S =& \varepsilon^7(-2u^2+(1+\alpha)u-7-3\alpha) \\ & \times(-2u^2+(1+\alpha)u-2)(2u^2+(4+2\alpha)u+7+3\alpha)^2(u+1)^3 \\ & \times\{(-4u^3+(3+\alpha)u^2-2(1+\alpha)u+6+2\alpha) \pm 2(-\zeta^3+3\zeta^2+2\zeta+1)u^2\}. \end{aligned}$$

The quantities inside the brackets are, respectively,

$$\begin{aligned} (-4u^3+(3+\alpha)u^2-2(1+\alpha)u+6+2\alpha)+2(-\zeta^3+3\zeta^2+2\zeta+1)u^2 \\ = -4(u+\zeta)(u+\zeta^3)(u-(1+\zeta)^2), \end{aligned}$$

and

$$\begin{aligned} (-4u^3+(3+\alpha)u^2-2(1+\alpha)u+6+2\alpha)-2(-\zeta^3+3\zeta^2+2\zeta+1)u^2 \\ = -4(u+\zeta^2)(u+\zeta^4)(u-\zeta^3(1+\zeta)^2). \end{aligned}$$

On the other hand, the factors of the quantity A are

$$2u^2+(-1+\alpha)u+2 = 2(u+\zeta)(u+\zeta^4),$$

while

$$-2u^2+(1+\alpha)u-2 = -2(u+\zeta^2)(u+\zeta^3).$$

Now, using the factorizations

$$-2u^2+(1+\alpha)u-7-3\alpha=-2(u-(1+\zeta)^2)(u-\zeta^3(1+\zeta)^2),$$
$$2u^2+(4+2\alpha)u+7+3\alpha=2(u-\zeta(1+\zeta)^2)(u-\zeta^2(1+\zeta)^2),$$

we find the two expressions Y_1 and Y_2 stated in the theorem. These factorizations also yield the factorization of the numerator and denominator of X in (2.2). $\square$

Remarks. The theorem shows that the quantities X and Y_i factor in a similar way over $\mathbb{Q}(\zeta)$ to the way that the quantity b factors:

$$b=\varepsilon^5\frac{[u-(1+\zeta)^2][u-\zeta(1+\zeta)^2][u-\zeta^2(1+\zeta)^2][u-\zeta^3(1+\zeta)^2][u-\zeta^4(1+\zeta)^2]}{(u+\zeta)(u+\zeta^2)(u+\zeta^3)(u+\zeta^4)(u+1)}.$$

The expression for X may be written as

$$X=-\varepsilon^4\frac{(u-(1+\zeta)^2)}{u+1}\frac{(u-\zeta(1+\zeta)^2)}{u+\zeta}\frac{(u-\zeta^2(1+\zeta)^2)}{u+\zeta^2}\frac{(u-\zeta^3(1+\zeta)^2)}{u+\zeta^3}\frac{u+\zeta}{u+1},$$

and the Y_i may be written in a similar form. Thus, the coordinates of $P=(X,Y_i)$ are products of linear fractional expressions in u.

3 Checking the formulas.

The curve E_5 is isomorphic to the curve

$$E':\ Y'^2=X^3+\frac{b^2+6b+1}{4}X^2+\frac{b(b+1)}{2}X+\frac{b^2}{4}$$

by the substitution $Y'=Y+\frac{1}{2}(1+b)X+\frac{b}{2}$. From [8], Ex. 3.7 the doubling formula on E' is given by

$$X(2P)=\frac{X^4-(b^2+b)X^2-2b^2X-b^3}{4p(X)},\quad X=X(P),$$

with $p(X)=X^3+\frac{b^2+6b+1}{4}X^2+\frac{b(b+1)}{2}X+\frac{b^2}{4}$; and

$$Y'(2P)=\frac{N(X)}{16p(X)^2}Y'(P),$$

with

$$N(X) = 2X^6+(b^2+6b+1)X^5+(5b^2+5b)X^4+10b^2X^3+10b^3X^2+(b^5+5b^4)X+b^5.$$

Taking the expression for $X = X(P)$ from (2.2), we have

$$X(2P) = \frac{(-7+3\alpha)}{4}\frac{(-2u^2+(1+\alpha)u-3\alpha-7)(2u^2+(4+2\alpha)u+3\alpha+7)}{(-2u^2+(1-\alpha)u-2)(u+1)^2},$$

which only differs from (2.2) in the denominator, where α has been replaced by its conjugate $-\alpha$. Notice that the numerator in this formula for $X(2P)$ is

$$\begin{aligned}&(-7+3\alpha)(-2u^2+(1+\alpha)u-3\alpha-7)(2u^2+(4+2\alpha)u+3\alpha+7)\\ &= (28-12\alpha)u^4+(12-4\alpha)u^3+8u^2+(12+4\alpha)u+28+12\alpha.\end{aligned}$$

This expression is invariant (except for a factor of u^4) under the mapping $(\alpha \to -\alpha, u \to 1/u)$. From (2.1) we see that this mapping also leaves the quantity b invariant, and takes the denominator of X (a symmetric polynomial in u) to the denominator of $X(2P)$ divided by u^4. Hence, $X(2P)$ is the X-coordinate in Theorem 2.1 corresponding to the pair $(-\alpha, 1/u)$, and we may state the following.

Theorem 3.1. *If X is given by (2.2), the X-coordinate of the double of the point $P = (X, Y_i)$ on E_5 is obtained by applying the mapping $(\alpha \to -\alpha, u \to 1/u)$ to the expression (2.2) or $(\zeta \to \zeta^2, u \to 1/u)$ to (2.3).*

Since the mapping $(\alpha \to -\alpha, u \to 1/u)$ has order 2, it is clear that $X(4P) = X(P)$ for either of the points $P = (X, Y_i)$ in Theorem 2.1. Applying the map $\sigma = (\zeta \to \zeta^2, u \to 1/u)$ to the quantity Y_1 in Theorem 2.1 yields

$$Y_1^\sigma = \bar{\varepsilon}^7\frac{[1-(1+\zeta^2)^2u]^2[1-\zeta^2(1+\zeta^2)^2u]^2[1-\zeta^4(1+\zeta^2)^2u]^2[1-\zeta(1+\zeta^2)^2u]}{(1+\zeta^4u)^2(1+\zeta u)(1+\zeta^3u)(u+1)^3},$$

and therefore, since $\frac{1}{(1+\zeta^2)^2} = \zeta^2(1+\zeta)^2$ and

$$(\zeta^2+\zeta^3)^7\cdot(1+\zeta^2)^{14}\cdot\zeta = 21+13(\zeta^2+\zeta^3) = -\varepsilon^7,$$

we find that

$$Y_1^\sigma = \varepsilon^7\frac{[u-(1+\zeta)^2]^2[u-\zeta(1+\zeta)^2][u-\zeta^2(1+\zeta)^2]^2[u-\zeta^3(1+\zeta)^2]^2}{(u+\zeta)^2(u+\zeta^2)(u+\zeta^4)(u+1)^3}.$$

If $P = (X, Y_1)$, this gives an expression for $Y_1^\sigma = Y(\pm 2P)$.

Since $\sigma^2 = (\zeta \to \zeta^4, u \to u) = (\zeta \to \zeta^{-1}, u \to u)$, we also have

$$Y_1^{\sigma^2} = \varepsilon^7 \frac{[u-(1+\zeta)^2][u-\zeta(1+\zeta)^2]^2[u-\zeta^2(1+\zeta)^2]^2[u-\zeta^3(1+\zeta)^2]^2}{(u+\zeta)(u+\zeta^2)(u+\zeta^3)^2(u+1)^3},$$

which coincides with Y_2. We have therefore that

$$P^{\sigma^2} = (X, Y_1)^{\sigma^2} = (X, Y_2) = -P,$$

and Theorem 3.1 yields

$$P^\sigma = (X, Y_1)^\sigma = \pm 2P.$$

Since σ is an automorphism of the extension $\mathbb{Q}(\zeta, u)/\mathbb{Q}(b)$, this shows that

$$-P = (P^\sigma)^\sigma = \pm 2P^\sigma = 4P,$$

and verifies that $4P = -P$, i.e. $5P = O$.

4 Appendix.

The roots of the equation $g(X) = 0$ are found using Watson's method and the following quantities defined in [1]. First, if

$$a_1 := \frac{1}{20}(\alpha - 5)(-3 - \alpha - 7b + 3b\alpha - 2b^2)$$

is the coefficient of X^4 in $g(X)$, we have

$$f(x) = g(x - \frac{a_1}{5}) = x^5 + 10Cx^3 + 10Dx^2 + 5Ex + F,$$

where:

$$C = \frac{1}{4000}(-3+\alpha)(b+3+\alpha)(-4b+3+\alpha)(-2b-11+5\alpha)(2b+11+5\alpha),$$
$$D = \frac{-1}{100000}(-5+2\alpha)(-8b^3-27b^2+11\alpha b^2-41b-19\alpha b+4+4\alpha)$$
$$\times(-2b-11+5\alpha)(2b+11+5\alpha)^2,$$

$$E = \frac{1}{500000}(-7+3\alpha)(-6b^5-60b^4+6\alpha b^4-135b^3+47\alpha b^3+505b^2+229\alpha b^2$$
$$-150b-72\alpha b+12+6\alpha)(-2b-11+5\alpha)(2b+11+5\alpha)^2,$$
$$F = \frac{-1}{12500000}(-25+11\alpha)(-8b^7-133b^6+5\alpha b^6-707b^5+115\alpha b^5+3790b^4$$
$$+2900\alpha b^4-15405b^3-6475\alpha b^3+5326b^2+2400\alpha b^2-794b-360\alpha b+44+20\alpha)$$
$$\times(-2b-11+5\alpha)(2b+11+5\alpha)^2.$$

Incidentally, this shows that the polynomial $f(x)$ is irreducible over $\mathbb{Q}(\alpha, b)$, by an analogue of Eisenstein's theorem, because of the factor $-2b-11+5\alpha$ in each of the coefficients. This yields:

$$K = E+3C^2$$
$$= \frac{1}{8000}(-9+4\alpha)b^2(-2b+29+13\alpha)(-2b-11+5\alpha)(2b+11+5\alpha)^2;$$
$$L = -2DF+3E^2-2C^2E+8CD^2+15C^4$$
$$= \frac{1}{140800000}(-35+16\alpha)b^4(-2b+1+\alpha)(22b-19+13\alpha)(-2b-11+5\alpha)^2$$
$$\times(2b+11+5\alpha)^4;$$
$$M = CF^2-2DEF+E^3-2C^2DF-11C^2E^2+28CD^2E-16D^4+35C^4E$$
$$-40C^3D^2-25C^6$$
$$= \frac{1}{512000000}(-9+4\alpha)b^6(-2b^4-20b^3+2\alpha b^3-11b^2-11\alpha b^2-35b-13\alpha b+3+\alpha)$$
$$\times(-2b-11+5\alpha)^3(2b+11+5\alpha)^5.$$

The discriminant of $f(x)$ is

$$\delta = \frac{1}{1024000}(123-55\alpha)(-2b-11+5\alpha)^4(2b+11+5\alpha)^8b^{14},$$

so that

$$\sqrt{\delta} = \frac{1}{1600}\alpha\left(\frac{1-\alpha}{2}\right)^5(-2b-11+5\alpha)^2(2b+11+5\alpha)^4b^7.$$

The polynomial

$$h(x) = x^6-\frac{K}{5}x^4+\frac{L}{125}x^2-\frac{\alpha\sqrt{\delta}}{390625}x+\frac{M}{3125}$$

has the quantity

$$\theta = \frac{1}{50} b(b^2 + 11b - 1)$$

as a root. Treating b as an indeterminate, $\theta \neq 0, \pm C$. Hence, Theorem 1 in [1] applies. We take

$$T = \frac{1}{20000}(5-\alpha)b(b-2+\alpha)(-2b-11+5\alpha)(2b+11+5\alpha)^2$$

to be the solution of $p(T) = q(T) = 0$ in that theorem. Then (2.20) in [1] gives

$$\begin{aligned} R_1 &= \sqrt{(D-T)^2 + 4(C-\theta)^2(C+\theta)} \\ &= \frac{1}{4000}(3-\alpha)b(2b-1+\alpha)(-2b-11+5\alpha)(2b+11+5\alpha)^2, \end{aligned}$$

and

$$\begin{aligned} R_2 &= \frac{C(D^2-T^2) + (C^2-\theta^2)(C^2+3\theta^2-E)}{R_1\theta} \\ &= \frac{1}{2000}(-2+\alpha)b(-2b+1+\alpha)(-2b-11+5\alpha)(2b+11+5\alpha)^2. \end{aligned}$$

Next, we define the quantities

$$\begin{aligned} X' &= \frac{1}{2}(-D+T+R_1) = \frac{1}{2^6 5^5}(-5+2\alpha)(2b+1+\alpha)(-2b-11+5\alpha)^2(2b+11+5\alpha)^3, \\ Y &= \frac{1}{2}(-D-T+R_2) = \frac{1}{2^4 5^5}(-5+2\alpha)(-b+2+\alpha)^2(-2b-11+5\alpha)^2(2b+11+5\alpha)^2, \\ Z &= -C-\theta = \frac{1}{2000}(3-\alpha)(-b+2+\alpha)(2b+1+\alpha)(-2b-11+5\alpha)(2b+11+5\alpha). \end{aligned}$$

Then, according to Theorem 1 of [1], the quantity u_1 is the fifth root of the expression

$$\begin{aligned} u_1^5 &= \frac{X'^2 Y}{Z^2} = \frac{1}{2^{11}5^8}(-25+11\alpha)(-2b-11+5\alpha)^4(2b+11+5\alpha)^6 \\ &= \frac{1}{2^{10}5^5}\left(\frac{-5+\alpha}{10}\right)^5(-2b-11+5\alpha)^5(2b+11+5\alpha)^5 \\ &\quad \times \frac{2b+11+5\alpha}{-2b-11+5\alpha}. \end{aligned}$$

This shows that u_1 is a polynomial in b and α times u, where

$$u^5 = \frac{2b+11+5\alpha}{-2b-11+5\alpha},$$

as in Section 2; in fact, we have

$$\begin{aligned} u_1 &= \frac{-5+\alpha}{200}(-2b-11+5\alpha)(2b+11+5\alpha) \times u \\ &= \frac{5-\alpha}{100}(2b^2+22b-2)u = \frac{5-\alpha}{100}A_1 u. \end{aligned}$$

This gives the first degree term in u in the expression for the root X in Section 2. Similarly, the quantities

$$\begin{aligned} \bar{X} &= \frac{1}{2}(-D+T-R_1) \\ &= \frac{1}{100000}(-5+2\alpha)(-b+2+\alpha)(-2b-1+\alpha)^2(-2b-11+5\alpha)(2b+11+5\alpha)^2, \\ \bar{Y} &= \frac{1}{2}(-D-T-R_2) \\ &= \frac{1}{200000}(-5+2\alpha)(-2b-1+\alpha)(2b+1+\alpha)^2(-2b-11+5\alpha)(2b+11+5\alpha)^2, \\ \bar{Z} &= -C+\theta = \frac{1}{4000}(3-\alpha)(-2b-1+\alpha)(-2b-11+5\alpha)(2b+11+5\alpha)^2, \end{aligned}$$

yield the expressions

$$u_2 = \frac{\bar{X}}{\bar{Z}^2}u_1^2 = \frac{5-\alpha}{100}A_2u^2, \quad u_3 = \frac{\bar{X}\bar{Y}}{Z\bar{Z}^3}u_1^3 = \frac{5-\alpha}{100}A_3u^3,$$

$$u_4 = \frac{\bar{X}^2\bar{Y}}{Z^2\bar{Z}^4}u_1^4 = \frac{5-\alpha}{100}A_4u^4,$$

which are the second, third, and fourth degree terms in the expression for the root X, where

$$\begin{aligned} A_2 &= (b-2-\alpha)(-2b-11+5\alpha), \\ A_3 &= -\frac{1}{2}(2b+1+\alpha)(-2b-11+5\alpha), \\ A_4 &= -\frac{1}{2}(-2b-1+\alpha)(-2b-11+5\alpha). \end{aligned}$$

Together with the fact that $\frac{(5-\alpha)}{100}A_0 = -\frac{a_1}{5}$, this yields the expression for the root

$$X = u_1 + u_2 + u_3 + u_4 - \frac{a_1}{5} = \frac{(5-\alpha)}{100}(A_4u^4 + A_3u^3 + A_2u^2 + A_1u + A_0)$$

of $g(X) = 0$ in Section 2. By replacing u by $\zeta^i u$, for $0 \leq i \leq 4$, and solving the resulting system of linear equations for the powers of u, it is not hard to see that $\mathbb{Q}(\zeta, b, X) = \mathbb{Q}(\zeta, b, u)$ is the field generated over $\mathbb{Q}(\zeta, b)$ by the X-coordinates of the points of order 5 on E_5. This gives an alternate verification that u^5 is a Kummer element for the extension $\mathbb{Q}(\zeta, b, X)/\mathbb{Q}(\zeta, b)$, as in [9].

References

[1] M.J.Lavallee, B.K. Spearman, K.S. Williams, Watson's method of solving a quintic equation, JP J. Algebra, Number Theory & Appl. 5 (2005), 49-73.

[2] R. Lynch, Arithmetic on normal forms of elliptic curves, Ph.D. thesis, Purdue University, 2015.

[3] R. Lynch and P. Morton, The quartic Fermat equation in Hilbert class fields of imaginary quadratic fields, International J. of Number Theory 11(2015), 1961-2017.

[4] P. Morton, Explicit identities for invariants of elliptic curves, J. Number Theory 120 (2006), 234-271.

[5] P. Morton, The cubic Fermat equation and complex multiplication on the Deuring normal form, Ramanujan J. 25(2011), 247-275.

[6] P. Morton, Solutions of the cubic Fermat equation in ring class fields of imaginary quadratic fields (as periodic points of a 3-adic algebraic function), International J. of Number Theory 12(2016), 853-902.

[7] P. Morton, Solutions of diophantine equations as periodic points of p-adic algebraic functions, I, New York J. of Math. 22 (2016), 715-740.

[8] J.H. Silverman, *The Arithmetic of Elliptic Curves*, 2nd edition, Springer, 2009.

[9] H. Verdure, Lagrange resolvents and torsion of elliptic curves, Int. J. of Pure and Appl. Math. 33, No. 1 (2006), 75-92.

Dept. of Mathematical Sciences, LD 270
Indiana University - Purdue University at Indianapolis (IUPUI)
Indianapolis, IN 46202
e-mail: pmorton@iupui.edu